

ACCURATE PARAMETER ESTIMATION FOR COUPLED STOCHASTIC DYNAMICS

ROBERT AZENCOTT

Professor, Dept of Mathematics, University of Houston and Ecole Normale Sup, France

YUTHEEKA GADHYAN

PhD Student, Department of Mathematics, University of Houston, USA

ABSTRACT. We develop and implement an efficient algorithm to estimate the 5 parameters of Heston’s model from arbitrary given series of joint observations for the stock price and volatility. We consider the time interval T separating two observations to be unknown and estimate it from the data, thereby estimating 6 parameters with a clear gain in fit accuracy. We compare the maximum likelihood parameter estimates based on an Euler discretization scheme to analogous estimates derived from the more accurate Milstein discretization scheme; we derive explicit conditions under which the two set of estimates are asymptotically equivalent, and we compute the asymptotic distribution of the difference of the two set of estimates. We show that parameter estimates derived from the Euler scheme by constrained optimization of the approximate maximum likelihood are consistent, and we compute their asymptotic variances. Numerically, our estimation algorithms are easy to implement, and require only very moderate amounts of CPU. We have performed extensive simulations which show that for standard range of the process parameters, the empirical variances of our parameter estimates are correctly approximated by their theoretical asymptotic variances.

1. Introduction. Derivatives such as futures and options were introduced to be used as hedging tools, and are now heavily traded in the market. Trading prices for these derivatives are indicators of market expectations for the near future. Due to high volatilities, efficient hedging tools require accurate pricing of these derivatives. To evaluate option prices through robust model based inference from asset dynamics data, it is crucial that the underlying joint stochastic dynamics of the asset price and volatility be calibrated correctly. Our objective here is to propose a fast and robust parameter estimation algorithm for such pairs of coupled SDEs, with good quantitative control for the accuracy of these estimates.

Stock price volatilities are notoriously not constant in time [23]. Local volatility models consider the volatility to be dependent on time and the underlying asset [11][12][25]. Stochastic volatility models drive volatility and price dynamics by correlated Brownian processes [17][26][27][28][29]. Various methods have been proposed for estimation of stochastic volatility models [1][4][8][15][19]. [1] employs maximum likelihood using a Hermite approximation of the likelihood function. [4]

2000 *Mathematics Subject Classification.* Primary: 60H10, 60H35; Secondary: 62F12, 91B28.

Key words and phrases. Stochastic differential equations, parameter estimation, maximum likelihood estimators, Heston model, asset price dynamics, volatility dynamics.

The second author is supported by NSF Grant DMS-0412267, NSF Grant ATM-0417867 and The Methodist Hospital Research Institute (Cardiology).

develops a weighted non parametric approach to determine the risk-neutral measure of the future states of the market. Bayesian estimation is proposed in [19]. We focus our analysis on parameter estimation for the often used pair of coupled SDEs introduced by Heston to price options [17]. Feller’s “square root” SDE [13] for volatility was first applied by Cox, Ingersoll and Ross to model short term interest rates [9]. We do not study here the pragmatic suitability of such SDEs to model the joint dynamics of stock price and volatility, but focus on generating easily computable parameter estimates from finite sets of observed data, and on the simultaneous evaluation of the estimates’ accuracy. In this context, there is no closed form expression for the log-likelihood or the joint density of price and volatility.

Heston’s pair of coupled SDEs for price and volatility involve 5 unknown parameters. We derive robust parameter estimates from an approximate maximum likelihood principle applied to the Euler discretization scheme for SDEs. The time interval T between consecutive observations is not fixed a priori but also simultaneously estimated from the data.

We evaluate asymptotic variances for these consistent parameter estimates, and compare their accuracy to a second group of estimates based on the more elaborate Milstein discretization scheme for SDEs. The Milstein scheme approximates underlying diffusion processes with L_2 norm accuracy equivalent to T , while the Euler scheme accuracy is of the order of \sqrt{T} [22]. We derive explicit conditions to force the Euler scheme parameter estimates to remain very close to the Milstein scheme estimators, and we compute the asymptotic distribution of the differences between these two types of estimators. We show that the estimators obtained are consistent. Numerically, our estimation algorithms are easy to implement, and require only very moderate amounts of CPU. Our extensive simulations show that for standard ranges of the process parameters, the empirical variances of our parameter estimates are correctly approximated by their theoretical asymptotic variances.

2. Heston’s coupled SDEs for volatility and asset price. Consider Heston’s classical coupled SDE system (under market measure) [17]

$$dS_t = \left(\mu - \frac{1}{2}X_t\right)dt + \sqrt{X_t}dW_1(t) \tag{1}$$

$$dX_t = \kappa(\theta - X_t)dt + \gamma\sqrt{X_t}dW_2(t). \tag{2}$$

At time t , $Y_t = \exp(S_t)$ is the asset price and $\sqrt{X_t}$ is its volatility. The squared volatility X_t is a mean reverting square root process [9] with mean reversion speed $\kappa > 0$ and volatility γ . Processes $W_1(t)$, $W_2(t)$ are correlated Brownian motions with $E[dW_1(t)dW_2(t)] = \rho dt$. Parameters μ and θ are respectively the mean rate of return of the asset under market measure, and its long run average variance. Existence and uniqueness of the solution for these SDEs is well known[16]. Denote the vector of model parameters by $PAR = (\mu, \kappa, \theta, \gamma, \rho)$ and assume the following classical constraints to be satisfied

$$\kappa > 0, \quad \gamma > 0, \quad \theta > 0, \quad -1 \leq \rho \leq 1, \quad 2\kappa\theta > \gamma^2,$$

The last constraint $2\kappa\theta > \gamma^2$ is Feller’s square root condition [13] ensuring that X_t stays a.s. away from 0, so that $X_0 > 0$, then forces X_t to remain positive. Let

$$m = \frac{2\kappa\theta}{\gamma^2} \text{ and } \nu = \frac{\gamma^2}{2\kappa}.$$

The conditional density of X_t given X_s is a non central χ^2 with $2m$ degrees of freedom. Its steady state density $p(x)$ is given by $\frac{x^{m-1} \exp(\frac{-x}{\nu})}{\nu^m \Gamma(m)}$ [9].

By integrating 2 and taking expectations, one easily obtains explicit expressions for $E[X_t]$ and $E[X_t^2]$, which show that as $t \rightarrow \infty$ the mean and variance of X_t converge at exponential speed towards θ and $\frac{\gamma^2 \theta}{2\kappa}$ respectively.

The joint density of $\{S_t, X_t\}$, has no explicit closed form. We want to estimate the 5 model parameters given a finite series of joint observations for the stock price $\{Y_t\}$ and square of volatility $\{X_t\}$. Note that in practice the volatility X_t is not directly observed. Good practical estimates of X_t have been proposed and used [2][3][7][10][14]. To focus on parameter estimation, we will assume here that the volatility is directly observed.

Let $(U_0, U_1, \dots, U_{N+1})$ and $(V_0, V_1, \dots, V_{N+1})$ be $N + 2$ joint observations for the log of asset price and the square of volatility at time points $t_0 < t_1 < \dots < t_{N+1}$, so that

$$U_n = S_{t_n} \text{ and } V_n = X_{t_n},$$

We assume that the time interval $T = t_{n+1} - t_n$ is fixed but unknown, to inject an adjustable time scale in model fitting. We want to select optimal T and PAR to achieve the best fit of the data by Heston's SDE system.

3. Approximate log-likelihood based on Euler Discretization. Euler discretization scheme for the SDE system provides the following difference equations, linking the one step differences ΔU_n , ΔV_n , $\Delta W_1(n)$ and $\Delta W_2(n)$,

$$\begin{aligned} \Delta U_n &= (\mu - \frac{1}{2}V_n)T + \sqrt{V_n}\Delta W_1(n) \\ \Delta V_n &= \kappa(\theta - V_n)T + \gamma\sqrt{V_n}\Delta W_2(n). \end{aligned} \tag{3}$$

Due to the presence of a square root term, the coefficients of our SDEs do not satisfy a global Lipschitz condition and therefore standard results on L_2 convergence for the Euler discretization do not apply here. However one can show that, under the parametric restrictions stated above, if the process and its Euler discretized approximation are both killed at the first (random) time the volatility becomes smaller than a fixed $\epsilon > 0$, then weak convergence of the Euler approximation scheme will hold in the associated space of continuous paths, endowed with an adequate natural metric. In particular, in order to ensure that V_n remains in $(0, \infty)$, sufficiently small discretization step sizes have to be imposed. The recent paper [20] develops efficient modified Ito-Taylor schemes to overcome both these difficulties, but we have not yet implemented their interesting approximation scheme. From 3 we get

$$\begin{aligned} \sqrt{V(n)}\Delta W_1(n) &= \Delta U(n) - (\mu - \frac{1}{2}V(n))T \\ \gamma\sqrt{V(n)}\Delta W_2(n) &= \Delta V(n) - \kappa(\theta - V(n))T, \end{aligned}$$

The independent random vectors $Z_{n+1} = (\Delta W_1(n), \gamma\Delta W_2(n))$ are Gaussian with

zero mean, variances T and $\gamma^2 T$, and covariance $\rho\gamma T$. Denote the coordinates of Z_{n+1} by $(z_1(n+1), z_2(n+1))$. The joint density of $\{Z_1\}, \dots, \{Z_N\}$ is:

$$f(Z_1, Z_2, \dots, Z_N) = \prod_{n=1}^N f(Z_n).$$

The maximum likelihood principle leads us to select the 6 unknown parameters T and PAR to maximize the log-likelihood LL given by

$$LL =: \ln f(Z_1, Z_2, \dots, Z_N) = \sum_{n=1}^N \ln f(Z_n).$$

The bivariate normal density of Z_n is completely determined by the covariance matrix of Z_n , and an easy computation then yields

$$\begin{aligned} \frac{1}{N} LL = & -\frac{1}{2} \ln(2\pi\gamma^2 T(1-\rho^2)) - \\ & \frac{1}{2N\gamma^2 T(1-\rho^2)} \sum_{n=1}^N (\gamma^2 z_1^2(n) - 2\rho\gamma z_1(n)z_2(n) + z_2^2(n)), \end{aligned} \quad (4)$$

$$\text{where } z_1(n) = \frac{\Delta U_n - (\mu - \frac{1}{2}V_n)T}{\sqrt{V_n}} \text{ and } z_2(n) = \frac{\Delta V_n - \kappa(\theta - V_n)T}{\sqrt{V_n}}.$$

A necessary condition to maximize LL is to let all its first derivatives be equal to zero :

$$\frac{\partial L}{\partial \theta} = \sum_{n=1}^N ((z_2(n) - \rho\gamma z_1(n)) \frac{\partial z_2(n)}{\partial \theta}) = 0, \quad (5)$$

$$\frac{\partial L}{\partial \kappa} = \sum_{n=1}^N ((z_2(n) - \rho\gamma z_1(n)) \frac{\partial z_2(n)}{\partial \kappa}) = 0, \quad (6)$$

$$\frac{\partial L}{\partial \gamma} = NT\gamma^2(1-\rho^2) + \rho\gamma \sum_{n=1}^N z_1(n)z_2(n) - \sum_{n=1}^N z_2^2(n) = 0, \quad (7)$$

$$\begin{aligned} \frac{\partial L}{\partial \rho} = & N\rho(1-\rho^2)\gamma^2 T + (\gamma + \rho^2\gamma) \sum_{n=1}^N z_1(n)z_2(n) \\ & - \rho \sum_{n=1}^N (\gamma^2 z_1^2 + z_2^2) = 0, \end{aligned} \quad (8)$$

where $z_1(n), z_2(n)$ are as in equation 4 and $L = -LL$.

For the derivative $\frac{\partial L}{\partial \mu}$, we use the simpler log-likelihood based only on the SDE verified by S_t , which does not make a significant difference in the estimates:

$$\frac{\partial L}{\partial \mu} = \sum_{n=1}^N \left(\frac{\Delta U_i - (\mu - 1/2V_n)T}{V_n} \right) = 0. \quad (9)$$

Solving the 5 equations 5-9 is achieved numerically by gradient descent, in order to compute the maximum likelihood parameter estimates based on Euler discretization. We come back to the detailed study of these estimates below. In principle a more accurate discretization scheme with faster speed of convergence to the true SDE solutions could also be used to generate an approximation of the log-likelihood, and then could provide parameter estimates by maximizing the log-likelihood. We first show why, for parameter estimation, there is no real advantage in using more

precise but more complex discretizations. We focus below on the Milstein discretization scheme, which approximates in L_2 the true SDE solution with accuracy of order T , instead of the \sqrt{T} accuracy provided by the Euler scheme[22][20].

4. **Milstein discretization scheme.** Equations 1 and 2 in integral form are:

$$\begin{aligned} S_t &= S_{t_o} + \int_{t_o}^t (\mu - \frac{1}{2}X(s))ds + \int_{t_o}^t \sqrt{X(s)}dW_1(s) \\ X_t &= X_{t_o} + \int_{t_o}^t \kappa(\theta - X(s))ds + \int_{t_o}^t \gamma\sqrt{X(t)}dW_2(s). \end{aligned}$$

Expand the integrals by Stochastic Taylor expansion[6] to obtain the following discretization:

$$\begin{aligned} \Delta U_n &= (\mu - \frac{1}{2}V_n)T + \sqrt{V_n}\Delta W_1(n) + \frac{\gamma}{2}A_n \\ \Delta V_n &= \kappa(\theta - V_n)T + \gamma\sqrt{V_n}\Delta W_2(n) + \frac{\gamma^2}{2}B_n, \end{aligned} \tag{10}$$

where

$$A_n = \int_{t_n}^{t_{n+1}} \int_{t_n}^s dW_2(u)dW_1(s), \quad B_n = \int_{t_n}^{t_{n+1}} \int_{t_n}^s dW_2(u)dW_2(s)$$

are martingales with mean zero [24], variance $\frac{T^2}{2}$ and $Cov(A_n B_n) = \frac{\rho T^2}{2}$.

As seen above for the Euler scheme, an approximate log-likelihood can be formally computed from the Milstein difference equation, and then one can compute the first derivatives of this approximate log-likelihood to set them equal to zero. The derivatives of the additional terms in the Milstein scheme are zero with respect to all parameters except γ . The functional form of the first order equations for $\kappa, \theta, \gamma, \rho$ is hence essentially the same for both discretizations.

Denote this system of first order equations by $F(PAR, \mathbf{Z})$ where $PAR = (\kappa, \theta, \gamma, \rho)$, $\mathbf{Z} = (z_1(1), z_1(2), \dots, z_1(N), z_2(1), z_2(2), \dots, z_2(N))$. We do not need to consider the equation corresponding to μ here since conditions corresponding to μ can be easily analyzed separately. Denote by (PAR_E, \mathbf{Z}_E) and (PAR_M, \mathbf{Z}_M) the parameter estimates and data vectors respectively corresponding to the Euler and the Milstein schemes. Define

$$\Delta PAR = PAR_M - PAR_E \quad \text{and} \quad \Delta Z = \mathbf{Z}_M - \mathbf{Z}_E,$$

A first order Taylor expansion of F at the point (PAR_E, \mathbf{Z}_E) gives:

$$\begin{aligned} F(PAR_M, \mathbf{Z}_M) &= F(PAR_E, \mathbf{Z}_E) + \frac{\partial F}{\partial PAR} \cdot \Delta PAR + \\ &\quad \sum_{i=1}^N \left(\frac{\partial F}{\partial z_1(i)} \cdot \frac{\partial z_1(i)}{\partial PAR} + \frac{\partial F}{\partial z_2(i)} \cdot \frac{\partial z_2(i)}{\partial PAR} \right) \cdot \Delta PAR + \frac{\partial F}{\partial Z} \cdot \Delta Z, \end{aligned} \tag{11}$$

where ΔPAR is a 4×1 vector, and all partial derivatives are matrices of adequate dimensions, evaluated at (PAR_E, \mathbf{Z}_E) . By definition of PAR_M , the left hand side of the above equation is zero. So it follows,

$$- \left(\frac{\partial F}{\partial PAR} + \sum_{i=1}^N \left(\frac{\partial F}{\partial z_1(i)} \cdot \frac{\partial z_1(i)}{\partial PAR} + \frac{\partial F}{\partial z_2(i)} \cdot \frac{\partial z_2(i)}{\partial PAR} \right) \right) \cdot \Delta PAR = \frac{\partial F}{\partial Z} \cdot \Delta Z. \tag{12}$$

The difference ΔPAR between our two vector estimates can be computed by inverting the preceding linear system, and of course depends on T . Using this expression for ΔPAR , and for any fixed small tolerance level η , we can compute a corresponding explicit upper bound for T , compelling each coordinate of ΔPAR to have variance less than η .

We also derive the asymptotic distribution of ΔPAR for large N . As $N \rightarrow \infty$ the coefficient matrix of ΔPAR divided by N in the preceding linear system converges to the constant 4×4 symmetric matrix f , where all coefficients are zero except the following ones

$$f_{11} = \frac{T}{2\kappa(1-\rho^2)}, \quad f_{22} = \frac{\kappa^2 T}{\gamma^2(1-\rho^2)\theta}, \quad f_{33} = \frac{2-\rho^2}{\gamma^2(1-\rho^2)},$$

$$f_{43} = f_{34} = \frac{-\rho}{\gamma(1-\rho^2)}, \quad f_{44} = \frac{1+\rho^2}{(1-\rho^2)^2}.$$

Denote the right hand side of the linear system 12 above by $R = \frac{\partial F}{\partial Z} \cdot \Delta Z$. We show that as $N \rightarrow \infty$ the random vector $\frac{R}{\sqrt{N}}$ is asymptotically Gaussian with mean zero and a limit covariance matrix Σ which we compute explicitly.

The two last results just mentioned show that as $N \rightarrow \infty$, the difference $\sqrt{N}(PAR_M - PAR_E)$ between the two groups of parameter estimates deduced from the Milstein and Euler schemes becomes asymptotically Gaussian with mean zero and an explicitly computed covariance matrix. Using these results, we have derived, for fixed large N , an explicit bound τ on T which will force the easily computed Euler scheme parameter estimates to remain reasonably close to the more accurate (but far harder to compute) estimators based on Milstein's discretization.

5. Constrained parameter estimation based on Euler discretization. As just shown, the Euler discretization, combined with an adaptive computable upper bound τ on T , generates controllably accurate parameter estimators for the Heston SDEs. The constrained estimation problem is now reduced to the maximization of $LL(T, PAR)$ as a function of T and PAR under the following constraints:

$$\kappa > 0, \quad \gamma > 0, \quad \theta > 0, \quad 2\kappa\theta > \gamma^2, \quad -1 \leq \rho \leq 1, \quad T < \min\left\{\tau, \frac{2}{\kappa}\right\},$$

where $T < \frac{2}{\kappa}$ ensures convergence of the Euler scheme[18]. The constraint on T depends on the parameter vector PAR . But PAR itself is unknown and its estimator depends on T . So we implement a sequence of alternating estimations by gradient descent for PAR and T , iterating until the estimates stabilize, according to the following steps :

- Start with initial value $PAR_0 = (\mu \ \theta \ \kappa \ \gamma \ \rho)$
- k th iteration generates estimate PAR_k from PAR_{k-1} by 1-step gradient descent
- Compute Control Bound τ_k for T to get $\tau_k = g(\text{data}, PAR_k)$
- Compute Unconstrained Estimate of T denoted $T_{opt} = h(\text{data}, PAR_k)$
- Compute $T_{k+1} = \min\{T_{opt}, \tau_k\}$
- Compute $PAR_{k+1} = \arg \max_{PAR} LL(T_{k+1}, PAR)$
- Continue till the sequence of estimates PAR_k and T_k stabilize.

At each iteration, the expression for T_{opt} is an explicit closed form function of the data and PAR .

5.1. Estimation of the vector of parameters *PAR*. At each iteration for a fixed value of T we get the following maximum likelihood estimates:

$$\hat{\mu} = \frac{\sum_{n=1}^N \frac{\Delta U_n}{V_n} + \frac{NT}{2}}{T \sum_{n=1}^N \frac{1}{V_n}} \quad \hat{\theta} = \frac{\sum_{n=1}^N \frac{\Delta V_n}{V_n} + \kappa NT - \rho\gamma \sum_{n=1}^N \frac{z_1(n)}{\sqrt{V_n}}}{\kappa T \sum_{n=1}^N \frac{1}{V_n}},$$

$$\hat{\kappa} = \frac{\rho\gamma \sum_{n=1}^N z_1(n) \sqrt{V_n} - \sum_{n=1}^N \Delta V_n + Ng}{T \left(\sum_{n=1}^N V_n - \frac{N^2}{\sum_{n=1}^N \frac{1}{V_n}} \right)},$$

where $g = \frac{\sum_{n=1}^N \frac{\Delta V_n}{V_n} - \rho\gamma \sum_{n=1}^N \frac{z_1(n)}{\sqrt{V_n}}}{\sum_{n=1}^N \frac{1}{V_n}}$. The parameters ρ and γ are estimated numerically since there is no explicit form for them in terms of the sample only. All the estimates are evaluated at each iteration which is performed till the estimates stabilize. In the next subsection we show the consistency and asymptotic variance of the estimates.

5.2. Consistency of estimates. We show the consistency in L_2 norm of all 5 parameter estimators using the explicit expressions obtained for them[5][21]. Rewrite the parameter estimates as:

$$\hat{\mu} = \mu + \frac{\sum_{n=1}^N \frac{\Delta W_1(n)}{\sqrt{V_n}}}{T \sum_{n=1}^N \frac{1}{V_n}},$$

$$\hat{\theta} = \theta + \frac{\gamma \sum_{n=1}^N \frac{\Delta W_2(n)}{\sqrt{V_n}} - \rho\gamma \sum_{n=1}^N \frac{z_1(n)}{\sqrt{V_n}}}{\kappa T \sum_{n=1}^N \frac{1}{V_n}},$$

$$\hat{\gamma} = \frac{-\rho \sum_{n=1}^N z_1(n) z_2(n) + \sqrt{(\rho \sum_{n=1}^N z_1(n) z_2(n))^2 + 4NT(1-\rho^2) \sum_{i=1}^N z_2^2(n)}}{2NT(1-\rho^2)}.$$

Dividing the numerator and denominator by N in the above equations and observing the independence of $\{\frac{\Delta W_1(n)}{\sqrt{V_n}}\}_{1 \leq n \leq N}$ and the fact that

$$\frac{\sum_{n=1}^N z_1(n) z_2(n)}{N} \rightarrow \rho\gamma T, \quad \frac{\sum_{n=1}^N z_2^2(n)}{N} \rightarrow \gamma^2 T,$$

we have

$$\hat{\mu} \rightarrow \mu, \quad \hat{\theta} \rightarrow \theta, \quad \hat{\gamma} \rightarrow \gamma.$$

The consistency of $\hat{\kappa}$ can be shown by observing that as $N \rightarrow \infty$

$$\hat{\kappa} \rightarrow \frac{\kappa T \left(\theta - \frac{N}{\sum_{n=1}^N \frac{1}{V_n}} \right)}{\frac{T}{N} \left(\sum_{n=1}^N V_n - \frac{N^2}{\sum_{n=1}^N \frac{1}{V_n}} \right)},$$

From the ergodicity of the Markov chain V_n we obtain $\frac{\sum_{n=1}^N V_n}{N} \rightarrow \theta$, which implies the consistency of $\hat{\kappa}$. The consistency of $\hat{\rho}$ is proved similarly. Therefore all the parameter estimates derived from the Euler scheme are consistent.

5.3. **Asymptotic variance of the Euler scheme parameter estimators.** For large values of N , $\frac{1}{N} \sum_{i=1}^N \frac{1}{V_i}$ is close to $\frac{1}{\theta}$ and we obtain

$$\begin{aligned} \text{Var}(\hat{\mu}) &\simeq \frac{\theta}{NT}, \quad \text{Var}(\hat{\theta}) \simeq \frac{\theta\gamma^2(1-\rho^2)}{\kappa^2NT}, \\ \text{Var}(\hat{\kappa}) &\simeq \frac{\gamma^2\kappa\theta^3}{2(\theta^2-1)^2N}, \quad \text{Var}(\hat{\rho}) \simeq \frac{1+\rho^2}{N}, \\ \text{Var}(\hat{\gamma}) &\simeq \frac{(4(2-s)+2\rho^2(s-1))\gamma^2}{2N(1-\rho^2)^2}, \quad s = \sqrt{(2\rho^4-3\rho^2+4)}. \end{aligned}$$

6. **Results of empirical tests.** We have performed many numerical simulations of the preceding iterative computation of estimates, and we have always observed that the estimation scheme is actually *convergent*. We observe that even for moderate values of N the estimates are good. We present below the estimates and empirical variances of these estimators for different values of N . The tests were made on simulated diffusion processes with simulation step $\delta = 10^{-3}$, $(U_0, U_1, \dots, U_{N+1})$, $(V_0, V_1, \dots, V_{N+1})$ observed at t_0, t_1, \dots, t_{N+1} with the true value of $T = t_{n+1} - t_n = .005 = 5\delta$. The empirical variance and mean of our parameter estimates were computed over 500 simulated trajectories. The empirical variances were compared to the theoretical variances obtained above. We observe a good fit between empirical and theoretical variances Fig.(1) as well as between empirical and theoretical means Fig.(2).




FIGURE 1. Comparison of empirical and theoretical variance. Empirical variance computed over 500 trajectories. (dashed line - theoretical, solid line - empirical)




FIGURE 2. Mean of estimates computed for $N = 500$ to $N = 20000$ at increments of 500. Mean computed over 500 trajectories. (dashed line - true value, solid line - mean of estimate)

REFERENCES

- [1] Y. Aït-Sahalia and R. Kimmel, *Maximum likelihood estimation of stochastic volatility models*, Journal of Financial Economics, **83** (2007), 413–452.
- [2] M. Avellaneda, *Minimum entropy calibration of asset pricing models*, International Journal of Theoretical and Applied Finance, **1** 1998, 447–472.
- [3] M. Avellaneda, D. Boyer-Olson, J. Busca and P. Friz, *Reconstruction of volatility: Pricing index options by the steepest descent approximation*, Risk Magazine, October(2002).
- [4] M. Avellaneda, R. Buff, C. Friedman, N. Grandchamp, L. Kruk, and J. Newman, *Weighted Monte Carlo: a new technique for calibrating asset pricing models*, International Journal of Theoretical and Applied Finance, **4** (2001), 91–119.
- [5] R. Azencott, *Densité des diffusions en temps petit : développements asymptotiques (part I)*, Séminaire de Probabilités de Strasbourg, **18** (1984), 402–498.
- [6] R. Azencott, *Formule de Taylor stochastique et développement asymptotique d'intégrales de Feynman*, Séminaire de Probabilités de Strasbourg, **S16** (1982), 237–285.
- [7] S. BenHamida and R. Cont, *Recovering volatility from option prices by evolutionary optimization*, Journal of Computational Finance, **8** Summer(2005).

- [8] C. Broto and E. Ruiz, *Estimation methods for stochastic volatility models: a survey*, Journal of Economic Surveys, **18** (2004), 613–649.
- [9] J.C. Cox, J.E. Ingersoll and S.A. Ross, *A theory of the term structure of interest rates*, Econometrica, **53** (1985), 385–408.
- [10] S. Crépey, *Calibration of the local volatility in a generalized Black-Scholes model using Tikhonov regularization*, SIAM Journal on Mathematical Analysis, **34** (2003), 1183–1206.
- [11] E. Derman and I. Kani, *The volatilities smile and its implied tree*, Goldman Sachs Quantitative Strategies Research Notes, January(1994).
- [12] B. Dupire, *Pricing with a smile*, Risk, January(1994), 18–20.
- [13] W. Feller, *Two singular diffusion problems*, Annals of Mathematics, **54** (1951), 173–182.
- [14] J.-P. Fouque, G. Papanicolaou and R. Sircar, “Derivatives in Financial Markets with Stochastic Volatility”, Cambridge University Press, 2000.
- [15] J.-P. Fouque, G. Papanicolaou and R. Sircar, *Mean-reverting stochastic volatility*, International Journal of Theoretical and Applied Finance, **3** (2000), 101–142.
- [16] I.I. Gihman and A.V. Skorohod, “Stochastic Differential Equations”, Springer-Verlag, 1972.
- [17] S.L. Heston, *A closed-form solution for options with stochastic volatility with applications to bond and currency options*, The Review of Financial Studies, **6** (1993), 327–343.
- [18] D.J. Higham and X. Mao, *Convergence of Monte Carlo simulations involving the mean reverting square root process*, Journal of Computational Finance, **8** (2005), 35–61.
- [19] E. Jacquier, N. Polson and P. Rossi, *Bayesian analysis of stochastic volatility models with fat-tails and correlated errors*, Journal of Econometrics, **122** (2004), 185–212.
- [20] A. Jentzen, P. E. Kloeden and A. Neuenkirch, *Pathwise approximation of stochastic differential equations on domains: higher order convergence rates without global Lipschitz coefficients*, Numerische Mathematik, **112** (2009), 41–64.
- [21] M.G. Kendall and A. Stuart, “Advanced Theory of Statistics”, Griffin, London, 1948.
- [22] P.E. Kloeden and E. Platen, “Numerical Solution of Stochastic Differential Equations”, Springer-Verlag, 1999.
- [23] J. Macbeth and L. Merville, *An empirical examination of the Black-Scholes call option pricing model*, The Journal of Finance, **34** (1979), 1173–1186.
- [24] P. Protter, “Stochastic Integration and Differential Equations”, Springer-Verlag, 2004.
- [25] M. Rubinstein, *Implied binomial trees*, Journal of Finance, **49** (1994), 771–818.
- [26] L. Scott, *Option pricing when the variance changes randomly: theory, estimation, and an application*, Journal of Financial and Quantitative Analysis, **22** (1987), 419–438.
- [27] E. Stein and J. Stein, *Stock price distributions with stochastic volatility: an analytic approach*, Review of Financial Studies, **4** (1991), 727–752.
- [28] J. Wiggins, *Option values under stochastic volatility*, Journal of Financial Economics, **19** (1987), 351–372.
- [29] Y. Zhu and M. Avellaneda, *A risk-neutral stochastic volatility model*, International Journal of Theoretical and Applied Finance, **1** (1998), 289–310.

Received August 2008; revised March 2009.

E-mail address: razencot@math.uh.edu

E-mail address: yutheeka@math.uh.edu